

広域環境モニタリングのための航空機を  
用いた放射性物質拡散状況調査

報告書

平成 24 年 6 月

独立行政法人 日本原子力研究開発機構

文部科学省の平成 23 年度放射能測定調査委託事業による委託業務として、日本原子力研究開発機構が実施した平成 23 年度「広域環境モニタリングのための航空機を用いた放射性物質拡散状況調査」の成果を取りまとめたものである。

## 目次

1. はじめに .....	1
2. 航空機モニタリングの経緯.....	2
3. 航空機モニタリングシステム.....	7
3.1 MEXT システム.....	7
3.2 NUSTEC システム .....	9
3.3 OYO システム .....	10
3.4 FUGRO システム .....	11
3.5 システムの比較.....	12
3.5.1 点線源によるレスポンス確認.....	12
3.5.2 リファレンスライン .....	14
3.6 計算コードによる MEXT システム特性評価.....	17
4. データ採取及び解析方法.....	19
4.1 事前確認.....	19
4.1.1 検出器の健全性確認.....	19
4.1.2 機体の汚染確認.....	19
4.1.3 ヘリコプタの機底の遮蔽.....	20
4.2 データ採取方法.....	22
4.2.1 ヘリコプタの選定.....	22
4.2.2 フライト方法.....	26
4.2.3 テストライン .....	27
4.2.4 結果の妥当性確認のための地上測定 .....	28
4.3 解析方法.....	29
4.3.1 線量換算係数(CD).....	30
4.3.2 空気減弱係数(AF).....	34
4.3.3 線量率-放射能換算係数.....	36
4.3.4 バックグラウンドの減算方法.....	37
4.3.4.1 BG-index.....	39
4.3.4.2 Cs-index.....	40
4.3.5 海拔高度による宇宙線の影響.....	41
4.3.6 Rn 子孫核種の影響.....	41
4.3.7 減衰補正.....	43
4.3.8 検出下限値及び信頼性.....	44
4.3.9 全線量率の検出下限.....	44
4.3.10 全線量率換算の不確かさ .....	45
4.3.11 放射性 Cs の沈着量の検出下限.....	47
4.3.12 放射性 Cs の沈着量の不確かさ .....	47
4.4 マッピング方法.....	48

4.4.1	補間方法.....	48
4.4.2	他の補間法との比較.....	49
5.	結果.....	52
5.1	線量率マップ.....	52
5.2	放射性 Cs の沈着量マップ.....	52
6.	考察.....	57
6.1	地上の測定結果との比較.....	57
6.2	測定時期における比較.....	59
6.3	三次元マップ.....	62
6.4	海拔高度と放射性物質の分布.....	63
6.5	降雪の影響.....	65
6.6	地質図との比較.....	66
6.6.1	北海道.....	67
6.6.2	東北地方.....	70
6.6.3	関東地方.....	73
6.6.4	中部地方.....	75
6.6.5	近畿地方.....	77
6.6.6	中国地方.....	79
6.6.7	四国地方.....	82
6.6.8	九州地方.....	84
6.6.9	南西諸島.....	87
6.6.10	地質図との対比のまとめ.....	87
7.	まとめ.....	89
	謝辞.....	90
	参考文献.....	90
Appendix 1	航空機モニタリング検討委員会.....	92
Appendix 2	テストラインデータ.....	123
Appendix 3	BG-index 及び Cs-index.....	136
Appendix 4	In-situ Ge 測定データ.....	142
Appendix 5	県別の線量率及び放射性 Cs の沈着量マップ.....	146

## Contents

1. Introduction.....	1
2. Progress of AMS.....	2
3. AMS system.....	7
3.1 MEXT system .....	7
3.2 NUSTEC system .....	9
3.3 OYO system .....	10
3.4 FUGRO system .....	11
3.5 Comparison of system.....	12
3.5.1 Response of system by the point source.....	12
3.5.2 Reference line.....	14
3.6 Radiological characterization of MEXT system by the calculation code.....	17
4. Data acquisition and analysis method.....	19
4.1 Preliminary survey .....	19
4.1.1 Detector adjustment.....	19
4.1.2 Self contamination of the helicopter.....	19
4.1.3 Shielding effect of the bottom of the helicopter .....	20
4.2 Data acquisition.....	22
4.2.1 Selection of the helicopter.....	22
4.2.2 Flight condition .....	26
4.2.3 Test line.....	27
4.2.4 Ground survey for validity of the AMS result.....	28
4.3 Analysis method.....	29
4.3.1 Conversion factor of dose rate (CD) .....	30
4.3.2 Attenuation factor (AF).....	34
4.3.3 Conversion factor from dose rate to radiocesium deposition .....	36
4.3.4 Discrimination of natural nuclide.....	37
4.3.4.1 BG-index.....	39
4.3.4.2 Cs-index.....	40
4.3.5 Influence of cosmic rays by the height above the sea level altitude.....	41
4.3.6 Influence of the Rn progeny .....	41
4.3.7 Decay correction .....	43
4.3.8 Detection limit and reliability.....	44
4.3.9 Detection limit of dose rate .....	44
4.3.10 Uncertainty of dose rate conversion.....	45
4.3.11 Detection limit of radiocesium deposition .....	47
4.3.12 Uncertainty of radiocesium deposition.....	47
4.4 Mapping method .....	48

4.4.1	Interpolation method .....	48
4.4.2	Comparison with other interpolation methods .....	49
5.	Result .....	52
5.1	Map of dose rate .....	52
5.2	Map of radiocesium dosition .....	52
6.	Consideration .....	57
6.1	Comparison with the ground survey.....	57
6.2	Comparison in the measurement date.....	59
6.3	3D map .....	62
6.4	Height above the sea level altitude and radiological distribution.....	63
6.5	Influence of the snow .....	65
6.6	Comparison with the geologic map .....	66
6.6.1	Hokkaido .....	67
6.6.2	Tohoku .....	70
6.6.3	Kanto .....	73
6.6.4	Chubu .....	75
6.6.5	Kinki.....	77
6.6.6	Chugoku .....	79
6.6.7	Shikoku .....	82
6.6.8	Kyusyu .....	84
6.6.9	Nansei-shoto.....	87
6.6.10	Summary of comparison with the geologic map.....	87
7.	Summary .....	89
	Acknowledgment .....	90
	Reference .....	90
Appendix 1	Committee of AMS .....	92
Appendix 2	Data of test-line .....	123
Appendix 3	BG-index and Cs-index .....	136
Appendix 4	Data of In-situ Ge .....	142
Appendix 5	Map of the dose rate and the radiocesium at each prefecture.....	146

### Table List

Table 2-1	Aerial monitoring schedule .....	3
Table 2-2	Detail schedule of Aerial monitoring.....	4
Table 3-1	Specification of 4 ARMS.....	7
Table 3-2	Response of ARMS by <sup>137</sup> Cs point source.....	12
Table 3-3	Response of ARMS at reference line.....	14
Table 3-4	Ratio of Peak and Compton of ARMS at reference line.....	16
Table 4-1	Shielding factor of helicopter .....	21
Table 4-2	Photo of helicopter .....	23
Table 4-3	Summary of CD and AF.....	31
Table 4-4	CD and AF in east Japan .....	32
Table 4-5	CD and AF in west Japan .....	33
Table 4-6	Limit of detection of ARMS.....	45
Table 4-7	Dispersion of CD and AF .....	46
Table 4-8	Number of measurement point .....	51
Table 6-1	Analysis condition of in ambient Fukushima Dai-ichi NPP.....	60

## Figure List

Fig. 3-1	Block diagram of MEXT system.....	8
Fig. 3-2	Block diagram of NUSTEC system .....	9
Fig. 3-3	Block diagram of OYO system .....	10
Fig. 3-4	Block diagram of FUGRO system .....	11
Fig. 3-5	Counting loss of NUSTEC and MEXT-1 system .....	13
Fig. 3-6	Spectrum of MEXT-1 using $^{137}\text{Cs}$ source.....	13
Fig. 3-7	Distance between detectors and $^{137}\text{Cs}$ source (MEXT-1) .....	13
Fig. 3-8	Distance between detectors and $^{137}\text{Cs}$ source (MEXT-2) .....	14
Fig. 3-9	Distance between detectors and $^{137}\text{Cs}$ source (MEXT-3) .....	14
Fig. 3-10	Sukagawa reference line.....	15
Fig. 3-11	Utsunomiya reference line.....	15
Fig. 3-12	Count rate of Sukagawa reference line.....	15
Fig. 3-13	Count rate of Utsunomiya reference line.....	15
Fig. 3-14	Gamma spectra of ARMS on the reference line (Sukagawa) .....	15
Fig. 3-15	Model of MEXT system.....	17
Fig. 3-16	Simulated spectrum of $^{137}\text{Cs}$ point source .....	18
Fig. 3-17	Energy response of MEXT system using EGS5 .....	18
Fig. 3-18	Distribution of angular response of MEXT system using EGS5.....	18
Fig. 3-19	Ratio of count rate at 300 m above the ground.....	18
Fig. 4-1	System check of MEXT-1 .....	19
Fig. 4-2	Decontamination of helicopter .....	20
Fig. 4-3	Shielding by fuel tank in BK117 .....	20
Fig. 4-4	Angler response of MEXT-1 in Helicopter .....	20
Fig. 4-5	Angler response of MEXT (Bell412) .....	21
Fig. 4-6	Angler response of MEXT (Bell430, S76).....	21
Fig. 4-7	Example of flight plan.....	26
Fig. 4-8	Image of test-line flight .....	27
Fig. 4-9	Dose rate of test-line.....	28
Fig. 4-10	Analysis flow of ARMS in west japan .....	29
Fig. 4-11	Analysis flow of ARMS in east japan .....	30
Fig. 4-12	Simulated attenuation factor using EGS5 (Source radius: 1000 m) .....	34
Fig. 4-13	Example for attenuation factor by hovering .....	35
Fig. 4-14	Example for attenuation factor .....	35
Fig. 4-15	Spectrum area for BG-index and Cs-index.....	38
Fig. 4-16	Background dose rate using NaI survey meter (2005-2009) <sup>12)</sup> .....	39
Fig. 4-17	Histogram of BG-index at MEXT-2.....	39
Fig. 4-18	Comparison with all count rate and >1,400 keV count rate at MEXT-2 .....	39



Fig. 4-19	Histogram of BG-index at MEXT-3 .....	40
Fig. 4-20	Histogram of BG-index at OYO.....	40
Fig. 4-21	Example for Cs-index in MEXT-1 .....	40
Fig. 4-22	Example for histogram of Cs-index in MEXT-1 .....	41
Fig. 4-23	Count rate of cosmic-ray above sea.....	41
Fig. 4-24	Count rate of Rn-progeny above sea .....	42
Fig. 4-25	CF in $\beta$ ( $\text{g}/\text{cm}^2$ ) <sup>8)</sup> .....	43
Fig. 4-26	<sup>134</sup> Cs/ <sup>137</sup> Cs and A .....	43
Fig. 4-27	Value of BG <sub>self</sub> each prefecture.....	45
Fig. 4-28	Relationship of AF and altitude.....	46
Fig. 4-29	Detection limit of radiocesium count rate .....	47
Fig. 4-30	Detection limit of radiocesium deposition ( $\text{kBq}/\text{m}^2$ ).....	47
Fig. 4-31	Methods of mapping.....	48
Fig. 4-32	Parameter of IDW method.....	49
Fig. 4-33	Comparison of map by approximate interpolation method .....	50
Fig. 5-1	Dose rate of Japan (This result includes the influence of natural radioactive nuclides.)... 53	
Fig. 5-2	Deposition of <sup>134</sup> Cs in Japan.....	54
Fig. 5-3	Deposition of <sup>137</sup> Cs in Japan.....	55
Fig. 5-4	Deposition of <sup>134</sup> Cs+ <sup>137</sup> Cs in Japan .....	56
Fig. 6-1	Comparison of ARMS and ground survey .....	57
Fig. 6-2	Comparison of ARMS and In-situ Ge measurement .....	58
Fig. 6-3	Comparison of summer 2011 and spring 2012 in ambient Fukushima Dai-ichi NPP .....	59
Fig. 6-4	Histogram of summer 2011 and spring 2012 in North Ibaraki.....	60
Fig. 6-5	Comparison of 4 <sup>th</sup> /3 <sup>rd</sup> in Fukushima (inside 80km from Fukushima Dai-ichi NPP) .....	61
Fig. 6-6	Histogram of 3 <sup>rd</sup> /4 <sup>th</sup> in Fukushima (inside 80km from Fukushima Dai-ichi NPP).....	61
Fig. 6-7	Examples of 3D map .....	62
Fig. 6-8	Distribution of radiocesium inventory each altitude 20 m .....	63
Fig. 6-9	Distribution map of the radiocesium at an altitude above sea .....	64
Fig. 6-10	Ratio map of 2012 / 2011 results in ambient Fukushima Dai-ichi NPP .....	65
Fig. 6-11	Contrast with a geologic map (Hokkaido).....	69
Fig. 6-12	Contrast with a geologic map (Tohoku) .....	72
Fig. 6-13	Contrast with a geologic map (Kanto).....	74
Fig. 6-14	Contrast with a geologic map (Chubu).....	76
Fig. 6-15	Contrast with a geologic map (Kinki) .....	78
Fig. 6-16	Contrast with a geologic map (Chugoku).....	81
Fig. 6-17	Contrast with a geologic map (Shikoku) .....	83
Fig. 6-18	Contrast with a geologic map (Kyusyu).....	86
Fig. 6-19	Contrast with a geologic map (Nansei-Shotou).....	88